Mem. S.A.It. Vol. 74, 607

Memorie della

Restoring full-disk images for atmospheric and instrumental degradation effects

S. Criscuoli[1] and I. Ermolli[1]

INAF Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monte Porzio Catone, Italia
e-mail: serena@mporzio.astro.it

Abstract. A numerical technique developed to recover atmospheric and instrumental degradation effects on full-disk images of the solar atmosphere acquired with a medium resolution will be presented. This technique, based on the method proposed by Walton and Preminger (1999), allows simultaneous determination of the undisturbed solar limb darkening profile and the characterization of the Point Spread Function in terms of a small number of analytic parameters. The application of the technique allows to remove stray light effects from images, to a great extent, while preserves the data photometry. The results obtained applying this technique to a sample of full-disk images taken with the PSPT telescopes will be summarized.

Key words. Techniques: Image processing – Sun: Fundamental parameters

1. Introduction

Ground-based images of the Sun are, in general, degraded by "stray light". This term refers to scattering and blurring by both the Earth's atmosphere and the telescope itself. Detailed studies of solar features such as sunspots, faculae and the network necessary to model solar irradiance variations, as well as measurements of the solar image geometry, require correction for the stray light. In the following we present a technique developed to remove stray light effects to the greatest extent possible from the full-disk images obtained with the Rome-PSPT.
The technique is founded on a Fourier transform-based image restoring algorithm early proposed by Toner and Jefferies (1993) and Toner, Jefferies and Duvall (1997), then modified especially for what concerns the PSF description procedures by Walton and Preminger (1999). With respect to the one presented by the latter authors, our technique differs in the PSF fitting performed in the frequency domain, rather than in the real space.
We investigated the effects of applying the technique on the photometry of the real data acquired by applying it first to simulated images of the solar photosphere. The

Send offprint requests to: S. Criscuoli
Correspondence to: INAF OAR, Via Frascati 33, 00040 Monte Porzio Catone

application of the developed technique results in good recovery of the underlying image geometry and photometry. We then evaluated the mean results obtained restoring real full-disk solar images obtained with the two PSPT telescopes operative at the Rome and the M. Loa sites.

2. Assumptions and data description

The image formation process at a certain time t, under the hypothesis of isoplanatism, is described in the frequency domain as follows:

$$I(u,v) = I_0(u,v) \cdot S(u,v) + N(u,v)$$

where I is the Fourier transform of collected image, $I_0(x,y)$ is the Fourier transform of the ideal not aberrated image, $S(u,v)$ is the OTF that describes both atmospheric and instrumental degradation, N is the noise and (u,v) are spatial frequencies.
The aim of the algorithm is to find a proper description for the OTF in order to deconvolve the collected image with a restaurative non parametric Optimum filter (Brault and White, 1971):

$$W(u,v) = \frac{|S(u,v)|^2}{|S(u,v)|^2 + |N(u,v)|^2}$$

The technique has been developed for application to 1kx1k images extracted from the archive of the daily observations carried out with the PSPT at the Rome Observatory (Ermolli et al. 1997, Centrone et al. 2001). These images correspond to the observations acquired each day at the three band-pass centered at CaIIK (393.3±0.25nm), Blue continuum (409.2±0.25nm) and Red continuum (607.1±0.5nm). PSPT images are obtained by a summation of many short exposure frames (30 msec) for a total integration time of about 1 second, that is much more than a typical atmospheric correlation time (10-20 msec). Moreover instrumental PSPT aberrations are estimated to be less than 1" so that we can assume isoplanatism and a radially symmetric PSF (Walton and Preminger, 1999); the latter is modeled as the sum of three gaussians, that take into account of the atmospheric and instrumental blurring, and a lorentian, that describes the atmospheric and instrumental scattered light:

$$s(r) = C_1 [Y] + Z$$

$$Y = C_2 e^{-(\frac{r}{b_1})^2} + C_3 e^{-(\frac{r}{b_2})^2} + C_4 e^{-(\frac{r}{b_3})^2}$$

$$C_1 = (1 - a_l); \quad C_2 = (1 - a_2)(1 - a_3)$$

$$C_3 = a_2(1 - a_3); \quad C_4 = a_3$$

$$Z = \frac{a_l}{A(r^2 + b_l^2)}$$

where r is the radial coordinate (distance from the center of the sun) and A is a normalization parameter. The PSF is so described by a proper choice of the set of coefficients $\alpha = (al, a2, a3, bl, b1, b2, b3)$.
The not aberrated center to limb intensity variation (CLV) can be modeled using 5th order polynomial expansions like proposed by Pierce and Slaughter (1977):

$$i(\lambda, r) = \Sigma_{n=0}^{n=N} a_n(\lambda) \mu^n(r)$$

where μ is the heliocentric angle and a_n are functions of the wavelength λ and are tabulated.
The experimental CLV $o(r)$ will then be described, for a fixed wavelength, as $o(r) = i(r) * s(r) + n(x,y)$.
This vector is derived by data interpolating on a 0.5 px regular grid, the vector which entries are the median value of constant area annuli centered at the solar disk center. This technique allows to reduce the asymmetries introduced by solar structures and also to increase the SNR, so that we can neglect the noise in the $o(r)$ expression.
The set of coefficients that describes the PSF is derived by a least square fit:

$$\chi^2 = \Sigma_i \frac{|O(u_i) - I(u_i) \cdot S(U_i)|^2}{\sigma_i^2}$$