



Gamma-ray bursts as tracers of star-formation rate and metallicity evolution with THESEUS

S. D. Vergani

GEPI, Observatoire de Paris, PSL Universit, CNRS, 5 Place Jules Janssen, 92190 Meudon, France, e-mail: susanna.vergani@obspm.fr

Abstract. The class of long gamma-ray bursts (LGRBs) is associated with the collapse of the most massive stars, making them a tool to investigate star formation in the early Universe. Furthermore, thanks to their exceptional brightness, LGRB afterglows can be used as extragalactic background sources capable of unveiling the properties of the cold/warm gas of their hosting galaxies, up to the highest redshifts. Therefore LGRBs allow, uniquely, the combination of the information on different phases of the gas, through the absorption lines present in the afterglow spectra and the emission (continuum and lines) properties obtained from host galaxy photometry and spectroscopy, once the afterglow disappeared.

To date, the main problem to carry out this kind of studies at very high redshift is the poor number of GRBs detected at $z > 5$. The *THESEUS* mission will hugely increase such number, as it is expected to detect > 100 GRBs at $z > 5$. Therefore, *THESEUS* will bring a unique and fundamental contribution to the understanding of the chemical enrichment of the universe and of the evolution of galaxies and star formation up to the highest redshift.

Key words. Gamma-ray burst: general – Galaxies: star formation – Galaxies: ISM – Galaxies: evolution – Galaxies: high-redshift – Galaxies: abundances – early Universe

1. Introduction

Long-duration GRBs (LGRBs) are produced by massive stars, and so track star formation, and in particular the populations of UV-bright stars responsible for the bulk of ionizing radiation production.

Indeed, different studies pointed out the possibility to use LGRBs as tracers of star-formation activity up to the highest redshifts (e.g.: Robertson & Ellis 2012; Kistler et al. 2013). The advantage of using LGRBs instead of galaxies is that GRB are detected through their gamma-ray emission and afterglows which are bright also at high redshift. On the other hand, most of past works suffer

of the fact that only 30% of GRBs has a measured redshift, and of the lack of statistics at high redshift.

In order to use LGRB as star-formation tracers and LGRB host galaxies as a representative population of star-forming galaxies, we need to understand the link between the LGRB rate and star formation rate, in the following referred to as LGRB efficiency. Theoretically, there are various reasons to expect a dependence of the LGRB efficiency on metallicity. Indeed, most theoretical models predict low-metallicity for the GRB progenitor star as a necessary condition to eliminate the progenitor star hydrogen envelope (which would smother the jet, and produce emission lines in the GRB-




Fig. 1. *Left:* Expected detection rate of GRBs, and uncertainty, of *THESEUS* compared to *Swift*. The improvement of the number of very high-redshift GRBs reflects in a much smaller error on the rates and therefore on the SFR density in function of redshift as shown in the right panel. *Right:* SFR density evolution, adapted from Robertson & Ellis (2012), showing the simulation of the information on the SFR density evolution that could be obtained thanks to the GRB detected by the *THESEUS* mission. *Credits:* *THESEUS* collaboration.

associated supernova that are not observed) without spinning down the angular momentum that the central engine needs to launch the jet.

Using the *Swift/BAT6* (Salvaterra et al. 2012) and SHOALS (Perley et al. 2016a) complete samples of LGRB host galaxies, respectively, Vergani et al. (2015, 2017), Japelj et al. (2016), and Perley et al. (2016b) showed that LGRBs prefer sub-solar metallicity host galaxies. This implies that, if metallicity is the only factor affecting the LGRB efficiency, LGRBs can be used to trace star formation rate (SFR) at high-redshift (see also Greiner et al. 2015) as at that epoch the bulk of star-forming galaxies fulfil the condition of low-metallicity required by LGRBs.

During four years of operations *THESEUS* will detect more than 100 LGRBs at $z > 5$, ~ 10 of which being at $z > 10$. The substantial improvement in the GRB detected at $z > 7 - 8$ will reduce the uncertainties on the rates and bring to tighter constraints on the SFR density in function of redshift (see Fig.1).

2. GRBs as tracers of metallicity evolution

Thanks to their exceptional brightness, only marginally decreasing with redshift, GRB af-

terglows can be used as powerful extragalactic background sources. The afterglow spectrum reveals the signatures of the gas associated with the GRB host, but also with the IGM and with other galaxies present along the GRB line of sight. The spectroscopy of LGRB optical afterglows provide a unique tool to unveil the properties of the ISM even in the faintest and most distant galaxies. The analysis of the absorption lines bring to a direct measurement of the column densities of the different elements present in the interstellar medium (ISM; neutral hydrogen, metals, molecules...). The most prominent absorption line seen in high-redshift GRB spectra is the Lyman- α line of neutral hydrogen (HI). From the fitting of the damped Lyman- α (DLA) profile, it is possible to obtain accurate measurement of the neutral hydrogen column density (NHI). The absorption lines of different metal species, both in their low- and high-ionization state, are also commonly detected. The ratio of the metal and NHI column densities give a direct determination of the metallicity of the gas. The metallicity and abundance measurements of the ISM of LGRB host galaxies can be used to constrain the chemical enrichment of star-forming galaxies in the Universe (see Fig. 2). Indeed, the association of LGRBs with massive stars

spectroscopy from the ground-based facilities, identifying the ISM absorption features even at the highest redshifts (Fig. 2). Thanks to these observations, it will be possible to study the chemical evolution of star-forming galaxies, in principle even beyond $z = 10$ (Fig. 2).

3. Conclusions

THESEUS will have a unique impact in the studies of the early universe and galaxy evolution, by hugely improving the GRB statistics at high redshift and identifying very-high redshift GRBs. Both factors are fundamental and will allow a high-impact use of GRBs as probes of the high-redshift Universe, determining the evolution of the star-formation rate density and metallicity up to the early Universe.

GRBs offer the unique opportunity to investigate systematically (and in principle at any redshift) both the cold/warm gas in the ISM (using the absorption lines seen in the afterglow spectra) and the ionised gas (using the emission lines of the host galaxy spectra). For high-redshift faint star-forming galaxies, the combination of the information obtained from both diagnostics will not be possible through *standard* galaxy studies, even with the *JWST*. Instead, LGRBs offer the possibility to reach this goal up to the highest redshift.

The ground- and space-based follow-up of the significant number of high-redshift GRBs detected by *THESEUS*, will provide a unique and detailed picture of the properties of the population of high-redshift galaxies.

References

- Greiner, J., Fox, D. B., Schady, P., et al. 2015, *ApJ*, 809, 76
- Japelj, J., Vergani, S. D., Salvaterra, R., et al. 2016, *A&A*, 590, A129
- Jonker, P., O'Brien, P., Amati, L., et al. 2013, [arXiv:1306.2336](https://arxiv.org/abs/1306.2336)
- Kistler, M. D., Yuksel, H., & Hopkins, A. M. 2013, [arXiv:1305.1630](https://arxiv.org/abs/1305.1630)
- Perley, D. A., Krühler, T., Schulze, S., et al. 2016a, *ApJ*, 817, 7
- Perley, D. A., Tanvir, N. R., Hjorth, J., et al. 2016b, *ApJ*, 817, 8
- Robertson, B. E. & Ellis, R. S. 2012, *ApJ*, 744, 95
- Salvaterra, R., Campana, S., Vergani, S. D., et al. 2012, *ApJ*, 749, 68
- Sparre, M., Hartoog, O. E., Krühler, T., et al. 2014, *ApJ*, 785, 150
- Vergani, S. D., Salvaterra, R., Japelj, J., et al. 2015, *A&A*, 581, A102
- Vergani, S. D., Palmerio, J., Salvaterra, R., et al. 2017, *A&A*, 599, A120